

A comment on the R^4 -coupling in (M)atrix Theory

Marco Serone

Dept. of Mathematics, University of Amsterdam

Plantage Muidergracht 24, 1018 TV Amsterdam, The Netherlands

e-mail: serone@wins.uva.nl

Abstract

By scaling arguments we show that the presence of a R^4 -term in the eleven dimensional supergravity effective lagrangian, if it is visible in (M)atrix theory, should produce a correction to the five-loop effective lagrangian of two moving D0-branes.

There is no doubt that during last year there has been an increasing interest about the conjecture of Banks, Fischler, Shenker and Susskind (BFSS) [1]. They basically consider a non-perturbative formulation of M-theory in a fixed kinematical regime as given by the effective Super Yang-Mills $U(N)$ quantum mechanics [2] governing the IR dynamics of a system of N D0-branes. More precisely the original prescription stated that the D0-brane two-derivative lagrangian, parametrized by the string coupling constant g_S and the string length l_S , is a non-perturbative description of M-theory compactified on a spatial circle of radius R and Planck length l_P with the identifications

$$R = g_S l_S, \quad l_P = g_S^{1/3} l_S \quad (1)$$

This correspondence is realized in the infinite momentum frame where for any state with positive momentum P along R ,

$$P = \frac{N}{R} \rightarrow \infty, \quad N \rightarrow \infty \quad (2)$$

Later on, Susskind [3] realized that Matrix Theory (MT) is meaningful even for N finite, if we consider it as M-theory on a *light-like* circle. In this way, he showed that MT at finite N is just the so-called Discrete Light Cone Quantization of M-theory. Up to now a lot of tests have been performed to the BFSS conjecture [4, 5, 6]¹, some of them performed at finite N [8, 9, 10], using Susskind's prescription. For the simplest case of compactifications on a circle, all of them showed an agreement between the MT and the Suga computation of same physical processes². The case of compactified MT, and in particular to non-trivial spaces, is more involved and present some problems [12].

Very recently Seiberg [13] (see also [14]) gave a prescription that helps to clarify the range of validity of MT and gives new evidence to the point of view of [3]. In few words, he derived

¹We cite here only those works that provided evidence of MT through some explicit scattering computations. See [7] and references therein for a recent and wider perspective of MT.

²Note, however, that very recently a discrepancy for multigraviton scattering [11] appeared.

Susskind’s conjecture, relating M-theory compactified on a light-like circle to another M-theory on a spatial circle. More precisely, his prescription implies that any amplitude $\mathcal{A}_{D0}(g_S, l_S)$ computed starting from the D0-brane lagrangian is related to the corresponding amplitude $\mathcal{A}_M(l_P, R)$ of the M-theory on the spatial circle of radius R and Planck length l_P in the following limits:

$$\lim_{g_S, l_S \rightarrow 0} \mathcal{A}_{D0}(g_S, l_S) = \lim_{R, l_P \rightarrow 0} \mathcal{A}_M(R, l_P) \quad (3)$$

with $l_S g_S^{-1/3} = \text{fixed}, \quad l_P^2 R^{-1} = \text{fixed}$

where N is again related to the momentum P as $P = N/R$. First of all, note that there is not anymore any N -limit in the correspondence, that is then valid for any (positive) value of N . The important point is that the limit on the left side of eq.(3) defines precisely the perturbative regime of MT, on the contrary of the original proposal, where an extrapolation at strong coupling of amplitudes was needed.

Despite its success, MT has not been up to now a source of predictions; it has been used instead to find in new ways results already known in string or field theory. MT, however, is supposed to be a non-perturbative formulation of M-theory and then, by definition, a microscopic theory underlying eleven dimensional supergravity. In this respect, it should remove the bad UV-divergences that plague 11D SUGRA, in much the same way string theory does. MT should be able, for instance, to fix the coefficient (otherwise divergent) multiplying (one of) the R^4 -term appearing in the 11D SUGRA action [15, 16]; even if this finite coefficient can be fixed by requiring consistency with string theory, an explicit MT result should be seen as an alternative and more direct procedure and, at the same time, as a strong check to the validity of the theory itself.

Aim of this note is then to show that, if this term is in anyway visible in MT, it should appear as a correction to the effective lagrangian of two moving D0-branes induced by a *non-planar diagram* at five loops.

We will see, moreover, that according to the correspondence (3), the N -finite MT before

the $g_S, l_S \rightarrow 0$ limits, can be seen as a “regularized version” of eleven dimensional supergravity (compactified on a circle); in this way, among the infinite series of terms appearing at any loop l in MT for the potential between two D0-branes, all the terms proportional to v^n , with $n > 2l + 2$, are simply an effect of the regularization and, after an appropriate rescaling, vanish in the limit (3). In this perspective, on the other hand the “would-be” 5-loop term, responsible in MT of the R^4 -coupling, diverges in the limit (3), reproducing the UV-divergence occurring in Suga for this term.

Let us then consider the effective lagrangian for two D0-branes moving with relative velocity v and separation r ; by dimensional analysis it is straightforward to see that at loop l , their effective lagrangian $L^{(l)}$ can be written as [17]:

$$L^{(l)} = g_S^{l-1} \sum_{n=0}^{\infty} C_{nl} v^2 \left(\frac{v^2}{r^4} \right)^n \left(\frac{1}{r^3} \right)^l l_S^{4n+3l-1} \quad (4)$$

where the powers of l_S are fixed requiring $L^{(l)}$ to have the dimension of a $(\text{length})^{-1}$ and we omitted the N -dependence that will be discussed later. It is known that at 1-loop, $L^{(1)} \sim v^4/r^7$ ($C_{01} = 0$), whereas the two-loop effective lagrangian starts as v^6/r^{14} [9] ($C_{02} = C_{12} = 0$). It has been conjectured by [10] that at l^{th} loop order, $C_{0l} = \dots = C_{l-1,l} = 0$, in order to reproduce classical long-distance supergravity potentials. We will see that from the supergravity point of view this assertion is consistent, provided that we neglect higher derivative operators in the low-energy effective action³. Consider indeed the interaction of two massless particles (in 11D), considered as external sources, with gravity. The corresponding action is simply ⁴:

$$S = -\frac{1}{2k_{11}^2} \int d^{11}x \sqrt{g} R + \int d^{11}x \sqrt{g} g_{\mu\nu} T^{\mu\nu} \quad (5)$$

where it is understood that one direction (the 11th) is compactified on a circle of radius

³This observation has been also independently done in a recent paper [18].

⁴For what follows we will need simply the Einstein term of the Suga action.

R_{11} , the gravitational constant $k_{11}^2 = 16\pi^5 l_P^9$ [5] and

$$T^{\mu\nu}(x) = \sum_{i=1,2} \frac{P_i^\mu P_i^\nu}{E_i} \delta^{(10)}(x^\mu - x_i^\mu(t)) \quad (6)$$

We choose

$$\begin{aligned} x_1^\mu(t) &= (t, \frac{v}{2}t, \frac{b}{2}, 0, \dots, 0, v_{11}t) \\ x_2^\mu(t) &= (t, -\frac{v}{2}t, -\frac{b}{2}, 0, \dots, 0, v_{11}t) \end{aligned} \quad (7)$$

with $v \ll v_{11} \sim 1$, with the corresponding momenta P_i^μ ($P^{11} = N/R_{11}$). According to eq.(3), it is meaningful to consider a loopwise expansion, with k_{11}^2 as Planck constant. Expand then eq.(5) around the flat metric $g_{\mu\nu} = \eta_{\mu\nu} + k_{11} h_{\mu\nu}$ and take the graviton propagator in the De Donder gauge:

$$\Delta^{\mu\nu,\rho\sigma}(x-y) = (\eta^{\mu\rho}\eta^{\nu\sigma} + \eta^{\mu\sigma}\eta^{\nu\rho} - \frac{2}{9}\eta^{\mu\nu}\eta^{\rho\sigma}) \frac{1}{2\pi R_{11}} \int \frac{d^{10}q}{(2\pi)^{10}} \frac{e^{iq(x-y)}}{q^2} \quad (8)$$

where we have fixed $q_{11} = 0$. It is then an easy exercise, using eqs.(6), (7) and (8), to obtain the familiar leading $v^4/(b^2 + v^2 t^2)^{7/2}$ dependence of the potential between the two sources: a simple algebra shows that the tensorial structure of eqs.(6) and (8) imply that each source carries effectively a power of $P_{11} \cdot v^2$, whereas the graviton propagator gives a factor $\sim 1/R_{11} r^7$; these are of course the leading terms in an expansion in $v \ll 1$. Changing to the string units g_S, l_S , we have exactly the same scaling provided by eq.(4). Note, however, that if we rescale both amplitudes and then take the limit (3), all the terms $C_{n1}, n > 1$ in eq.(4) vanish, as previously mentioned.

In this kind of expansion in k_{11} , all higher order corrections scale necessarily like $(k_{11}^2)^m \sim l_P^{9m}$, with m an integer. This condition is indeed satisfied by eq.(4), provided $l = n$. For instance, the next to leading correction to the potential is given by contracting three source terms with the term trilinear in $h_{\mu\nu}$, coming from the expansion of the scalar curvature R . Even if the full contribution is rather involved, it is easily seen that the end result goes as v^6/r^{14} . The next order, coming from the four-graviton term in R , scales like

$$V_R \sim \frac{l_P^{27} v^8 P_{11}^4}{R_{11}^3 r^{21}} \quad (9)$$

where the 9-dimensional integration over the position of the interaction vertex cancels a contribution of one of the four propagators and the two derivatives of the vertex itself, leading then to eq.(9). In MT this correction should be visible as the three-loop effect $l = n = 3$ in eq.(4). It is clear that these terms correspond to the classical expansion of the potential found in [9] with a source-probe analysis. It has been conjectured by [15] that the 11D SUGRA action should contain a R^4 -term, that is needed to reproduce some perturbative and non-perturbative terms in IIA,B string theory. For M-theory compactified on a circle R_{11} , the analysis of [15] shows the presence of two contributions for R^4 :

$$S_{R^4} = \frac{R_{11}}{k_{11}^{2/3}} \int d^{10}x \sqrt{g} t_8 t_8 R^4 \left(\frac{2\pi^2}{3} + 2\zeta(3) \frac{l_P^3}{R_{11}^3} \right) \quad (10)$$

up to an overall factor, where the tensorial structure $t_8 t_8 R^4$ is explicitly given in [19]. For the purpose of the present analysis, we will not need the detailed form of this term; from the SUGRA point of view, these couplings arise as counterterms coming from a one-loop four graviton scattering. The first term in eq.(10) is actually UV-divergent and its finite coefficient is fixed by consistency with IIA,B string theory and T-duality, whereas the second one is completely finite [16]. We will see that the second term in eq.(10) is not visible in our MT computation (it does not match with our scaling arguments); we will focus our attention in the following on the first term only.

The presence of the R^4 -coupling (10) to the effective SUGRA action produces a sub-leading correction to the effective potential of the sources; in particular its linear structure contains a four-graviton term that induces a correction proportional to v^8/r^{27} , where the additional six powers of r with respect to eq.(9) come from the six more derivatives contained in R^4 , compared to the four-graviton two-derivative term of the R -coupling. From eq.(4), we see that a term of this form in MT is possible for $l = 5, n = 3$ and would be an “anomaly” to the suggestion that $C_{0l} = \dots = C_{l-1,l} = 0$; this is however expected since the correction (10) presents a fractional power of the gravitational constant k_{11} . We will see that this is consistent with our scaling arguments; the correction induced by the (first) term in eq.(10)

is

$$V_{R^4}^{Sugra} = c_S l_P^{33} \frac{v^8}{r^{27}} \frac{P_{11}^4}{R_{11}^3} \quad (11)$$

where c_S is a numerical coefficient. The corresponding MT potential is

$$V_{R^4}^{MT} = c_M l_S^{26} g_S^4 \frac{v^8}{r^{27}} \quad (12)$$

Using eq.(1), we see that there is a complete matching between the scaling of both terms,

$$\frac{V_{R^4}^{Sugra}}{V_{R^4}^{MT}} = \frac{c_S}{c_M} \quad (13)$$

where we omitted the N -dependence in both eqs.(12) and (13), dependence that will be discussed in the following. The presence of a non-vanishing coefficient c_M in eq.(12) would be a clear signal that MT is in the right way. If we normalize to a finite value in MT the “standard” $v^{2(l+1)}$ -term, then the limit (3) gives a divergent result for $V_{R^4}^{MT}$, that indeed reproduces the UV-divergence of this term in Sugra. The g_S -dependence of eq.(12) and then the origin of five-loops is easily seen in string theory: it is known that a generic two-dimensional surface with g handles, c crosscaps and b boundaries has a string coupling constant dependence given by $g_S^{(2g-2+b+c)}$. The usual annulus computation ($g = c = 0, b = 2$) gives of course g_S^0 , whereas the two and three loop MT amplitudes responsible for corrections of order $v^6/r^{14}, v^8/r^{21}$ respectively, correspond to a sphere with 3,4 punctures ($g = c = 0, b = 3, 4$)⁵. The R^4 -coupling we have considered is just the term arising at one-loop in IIA string theory [15], as can be easily seen using again eq.(1); the corresponding surface is then a torus with four punctures ($g = 1, c = 0, b = 4$) giving a g_S^4 dependence, coming in MT precisely at five loops. Let us consider now the N -dependence of these amplitudes; the leading large N -behaviour of eq.(4) at loop l is given by N^{l+1} , that does not match at five loops with the Sugra result $\sim N^4$. This is indeed what we expect since we have just seen that the two-dimensional surface responsible of this interaction in string

⁵Each puncture corresponds, from the Sugra point of view, to an insertion of the source term given by eq.(5).

theory involves a genus one surface, that leads then to a *non-planar diagram* in the Super Yang-Mills quantum mechanics, and with a sub-leading N power-dependence. Since every boundary on the world-sheet involves a trace on the fundamental of the gauge group, each of them gives one power of N , that then matches precisely with the analysis above.

A comment on the 2^{nd} term of eq.(10) is however needed. As already mentioned, the correction induced by this term does not match with the MT parameters; it should be visible as a planar diagram at three-loops, according with the analysis above, since it arises at tree-level in IIA string theory; at this order, however, the only v^8 -term appearing is that induced by the four-graviton term of the scalar curvature R , as in eq.(9). This mismatching could be due to the fact that this term arises probably as a counterterm generated by integrating out Kaluza-Klein states with non-vanishing momentum around R_{11} and is then not visible in a MT computation where we do not allow exchange of momentum in the eleventh direction.

This discussion can be also extended to more general higher derivative terms; any non-vanishing term in, say, the effective potential of two moving D0-branes with $n < l$ in eq.(4), correspond to a counterterm of higher dimension in the SUGRA action (see also [18] for a similar remark). In this perspective MT could in principle predict the form of new couplings and the finite correct values of their coefficients, in general divergent in SUGRA, although they should come out from very laborious many-loops computations in the Super Yang-Mills quantum mechanics.

Finally, I would like to comment on other papers [17, 20], where it is shown that the presence of the R^4 -coupling in the supergravity effective action should give corrections to the potential of two gravitons going like v^8/r^{18} . In [17] it is argued that the N -dependence of this term is consistent with the two loop term ($l = 2, n = 3$) of eq.(4), whereas the analysis of [20] shows a disagreement in the powers of N , not reconcilable with MT⁶. The

⁶Note, however, that in a recent paper [21] a possible explanation to the N -scaling of [20] is given.

reason of the apparent conflict between the results of [17, 20] with those presented here, is due to the different approaches followed. As shown before, we treat the external states in sugra as classical spinless sources, corresponding to the MT background of two moving D0-branes. The limit we consider corresponds precisely to weakly coupled IIA string theory in 10 dimensions with two clusters of coincident D0-branes. On the other hand refs.[17, 20] consider the external states as pure (super-)gravitons, with definite spin. The v^8/r^{18} term above is indeed found by considering also graviton polarizations; this effect is then not visible in our approximation where the external states are spinless. On the other hand, the source-probe analysis of [17, 20] cannot capture the term considered in this paper, being a non-local term in the effective action for the probe⁷.

It is not excluded, of course, the simplest possibility that c_M in eq.(12) is zero, just for kinematical reasons. It has been shown in [9], for instance, that the gravitational field created by a massless graviton in the light-cone frame cannot receive any correction by possible R^n -terms present in the lagrangian, because they are all vanishing in this frame [22]. I do not exclude that this is indeed what happens; a more precise computation in Sugra should answer this question.

Needless to say, it would be of the utmost importance to understand how and where to find in MT compactified on a circle and on a torus the results of [15, 16] for R^4 in nine and eight dimensions. I believe that a clear understanding of these (and similar) higher derivative interactions will shed a new light on the range of validity of Matrix Theory as a non-perturbative formulation of M-theory.

Acknowledgements

It is a pleasure to thank R. Dijkgraaf, E. Verlinde and H. Verlinde for useful and enlightening discussions.

⁷I thank E. Keski-Vakkuri and P. Kraus for a correspondence that helped me to clarify these points.

References

- [1] T. Banks, W. Fischler, S. Shenker and L. Susskind, *M-theory as a matrix model: a conjecture*, Phys. Rev. **D55** (1997) 112, hep-th/9610043;
- [2] M. Claudson and M. Halpern, *Supersymmetric ground state wave functions*, Nucl. Phys. **B250** (1985) 689;
M. Baake, M. Reinicke and V. Rittenberg, *Fierz identities for real Clifford algebras and the number of supercharges*, J. Math. Phys. **26** (1985) 1070;
R. Flume, *On quantum mechanics with extended supersymmetry and non abelian gauge constraints*, Ann. Phys. **164** (1985) 189;
- [3] L. Susskind, *An other conjecture about (M)atrix Theory*, hep-th/9704080;
- [4] O. Aharony and M. Berkooz, *Membrane dynamics in (M)atrix Theory*, Nucl. Phys. **B491** (1997) 184, hep-th/9611215;
G. Lifschytz and S.D. Mathur, *Supersymmetry and membrane interactions in (M)atrix Theory*, hep-th/9612087;
G. Lifschytz, *Four-brane and six-brane interactions in (M)atrix Theory*, hep-th/9612223; *A note on the transverse five-brane in (M)atrix Theory*, hep-th/9703201;
I. Chepelev and A.A Tseytlin, *Long-distance interactions of D-brane bound states and longitudinal 5-brane in (M)atrix theory*, Phys. Rev. **D56** (1997) 3672, hep-th/9704127;
J.M. Pierre, *Interactions of eight-branes in string theory and (M)atrix Theory*, hep-th/9705110;
R. Gopakumar and S. Ramgoolam, *Scattering of zero branes off elementary strings in Matrix Theory*, hep-th/9708022;
- [5] D. Berenstein and R. Corrado, *M(atrix) Theory in various dimensions*, Phys. Lett. **B406** (1997) 37, hep-th/9702108;

- [6] J. Polchinski and P. Pouliot, *Membrane scattering with M-momentum transfer*, hep-th/9704029;
- [7] T. Banks, *Matrix Theory*, hep-th/9710231;
- [8] K. Becker and M. Becker, *A two loop test of (M)atrix Theory*, hep-th/9705091;
P. Kraus, *Spin-orbit interaction from Matrix Theory*, hep-th/9709199;
- [9] K. Becker, M. Becker, J. Polchinski and A.A. Tseytlin, *Higher order graviton scattering in (M)atrix Theory*, Phys. Rev. **D56** (1997) 3174, hep-th/9706072;
- [10] I. Chepelev and A.A. Tseytlin, *Long-distance interactions of branes: correspondence between supergravity and super Yang-Mills descriptions*, hep-th/9709087;
E. Keski-Vakkuri and P. Kraus, *Born-Infeld actions from Matrix Theory*, hep-th/9709122;
- [11] M. Dine and A. Rajaraman, *Multigraviton scattering in the Matrix Model*, hep-th/9710174;
- [12] M.R. Douglas, H. Ooguri and S.H. Shenker, *Issues in (M)atrix Theory compactification*, Phys. Lett. **B402** (1997), hep-th/9702203;
O.J. Ganor, R. Gopakumar and S. Ramgoolam, *Higher loop effects in M(atrix) orbifolds*, hep-th/9705188;
M.R. Douglas, *D-branes and Matrix Theory in curved space*, talk given at Strings '97, hep-th/9707228;
M.R. Douglas and H. Ooguri, *Why Matrix Theory is hard*, hep-th/9710178;
- [13] N. Seiberg, *Why is the Matrix Model correct ?*, hep-th/9710009;
- [14] A. Sen, *D0 branes on T^n and Matrix Theory*, hep-th/9709220;
- [15] M.B. Green and P. Vanhove, *D-instantons, strings and M-theory*, hep-th/9704145;

- [16] M.B. Green, M. Gutperle and P. Vanhove, *One loop in eleven dimensions*, hep-th/9706175;
- [17] P. Berglund and D. Minic, *A note on effective lagrangians in Matrix Theory*, hep-th/9708063;
- [18] S.P. de Alwis, *Matrix models and string world sheet duality*, hep-th/9710219;
- [19] A.A. Tseytlin, *Heterotic-type I superstring duality and low-energy effective actions*, Nucl. Phys. **B467** (1996) 383, hep-th/9512081;
- [20] E. Keski-Vakkuri and P. Kraus, *Short distance contributions to graviton-graviton scattering: matrix theory versus supergravity*, hep-th/9712013;
- [21] V. Balasubramanian, R. Gopakumar and F. Larsen, *Gauge theory, geometry and the large N limit*, hep-th/9712077;
- [22] G.T. Horowitz and A.R. Steif, *Spacetime singularities in string theory*, Phys. Rev. Lett. **64** (1990) 260.